



Mary Ann Liebert, Inc. publishers

Journals

Search

Alerts

## JOURNAL OF PALLIATIVE MEDICINE

[J Palliat Med.](#) 2018 Oct 1; 21(10): 1529–1530.

Published online 2018 Oct 9. doi: [10.1089/jpm.2018.0348](https://doi.org/10.1089/jpm.2018.0348)

PMCID: PMC6909735

PMID: [30312130](#)

### The Surprise Question as a Prognostic Tool #360

[Kate S. Jennings](#), MD, [Sean Marks](#), MD, and [Hillary D. Lum](#), MD, PhD

[Author information](#) [Copyright and License information](#) [Disclaimer](#)

[Go to:](#)

#### Background

---

CLINICIANS HAVE BEEN ENCOURAGED to utilize the surprise question (SQ)—“Would I be surprised if this patient died within 12 months?”—to identify patients at high one-year mortality risk. When clinicians answer “No—I would NOT be surprised if this patient died within 12 months,” the SQ may help clinicians identify patients with unmet palliative care needs who could benefit from advance care planning discussions and/or a palliative care referral.<sup>1</sup> This *Fast Fact* reviews the clinical utility of the SQ.

[Go to:](#)

#### Rationale of the SQ

---

As difficult as it is for clinicians to prognosticate accurately, multiple studies have shown that patients with incurable disease desire more prognostic information the sicker they get and prognosis is a major factor in preferences for rehospitalizations, life support, and cardiopulmonary resuscitation.<sup>2-4</sup> Although patients and surrogates often want *temporal prognostic predictions* (the clinician's estimated length of time he or she predicts the patient will live), clinicians are more accurate and willing to offer *probabilistic predictions* (the clinician's estimate, often in a percentage, of the chance of death in a set time frame, such as one year).<sup>4-6</sup> The SQ was designed as a clinical tool that generalist clinicians would utilize willingly and routinely to identify patients at risk of death in a year and thereby lead to more appropriate advance care planning, goals of care discussions, symptom management, and hospice referrals.

[Go to:](#)

## Effectiveness of the SQ

---

Although variations on the SQ have been described in the published literature, including “Would I be surprised if this patient died this hospitalization?” or “Would I be surprised if this patient died in 3 months?” the SQ most commonly referred is “Would I be surprised if this patient died within 12 months?” This SQ has been studied in diverse populations, including a general inpatient setting,<sup>7</sup> high-risk primary care clinic,<sup>8</sup> pediatric patients,<sup>9</sup> advanced kidney disease,<sup>10-12</sup> cancer,<sup>13,14</sup> acute surgical patients,<sup>15</sup> emergency department settings,<sup>1,16</sup> and nursing home settings.<sup>17</sup> In general, the SQ has performed modestly well in identifying patients with a prognosis of less than one year across these various patient populations.<sup>18,19</sup> Notable findings from these studies include:

- A meta-analysis of the SQ among 26 studies across these patient populations found that it had a pooled accuracy of 75%, a sensitivity of 67%, and specificity of 80%.<sup>18</sup>
- “Yes” answers appear to be much more accurate than “No” answers. The predictive value of a “Yes” answer was 93%, whereas only 37% for a “No” answer.<sup>18</sup> This means that the SQ is likely better designed to identify patients who will live more than a year versus the patients who live less than a year. It also suggests that there are “false positives” relatively common when clinicians answer “No” to the SQ.
- The SQ may be slightly more accurate for cancer patients (pooled accuracy 79%) and renal patients (76%) versus other disease groups (72%).<sup>18</sup> This may reflect a more predictable illness trajectory in cancer and renal disease. See *Fast Fact #326* for more information on illness trajectories.
- In most studies, the SQ was utilized as one aspect of a broad prognostic assessment that included clinician gestalt and/or other prognostic tools. Hence, used in isolation, its accuracy is unclear.<sup>18</sup>

[Go to:](#)

## Implementing the SQ Into Clinical Practice

The SQ can be helpful in identifying patients at risk of medical decline and death in a certain time frame when used as part of a larger prognostic assessment.<sup>19</sup> Yet, considering the relatively high false positive rate of a “No” answer, it is not established if the SQ is a cost-effective tool nor an effective way to trigger a palliative care consultation on its own. A consensus panel of experts suggested that a “No” answer triggers generalist clinicians to perform a primary palliative care assessment or screening for unmet palliative care needs.<sup>20</sup> Sentinel medical events such as hospitalization, decline in performance status, or

disease progression are reasonable triggers to implement the SQ into clinical practice. Note templates, medical rounding tools, and electronic medical record prompts are potential system-based approaches to accomplish this. Reasonable components of the primary palliative care assessment triggered by a “No” include<sup>20</sup>:

- Assessment for distressing physical, psychological, social, or spiritual concerns.
- Identification of whether the patient completed an advance directive such as a healthcare power of attorney that is available in the medical records.
- Assessment of patient, family, and/or surrogate's understanding of the underlying illness, treatment options, and prognostic trajectory.
- Assessment of decision-making capacity.
- Engagement in honest conversations about prognosis and medical expectations.
- Elicitation of the patient's care preferences and values.
- Consideration of whether a hospice referral would be appropriate.
- Consideration of whether a palliative care consultation may be beneficial.

[Go to:](#)

## References

---

1. Haydar SA, Almeder L, Michalakes L, et al.: Using the surprise question to identify those with unmet palliative care needs in emergency and inpatient settings: What do clinicians think. *J Palliat Med* 2017;20:729–735 [[PubMed](#)] [[Google Scholar](#)]
2. Weeks JC, Cook EF, O'Day SJ, et al.: Relationship between cancer patients' predictions of prognosis and their treatment preferences. *JAMA* 1998;279:1709–1714 [[PubMed](#)] [[Google Scholar](#)]

3. Murphy DJ, Burrows D, Santilli S, et al.: The influence of the probability of survival on patients' preferences regarding cardiopulmonary resuscitation. *NEJM* 1994;330:545–549 [[PubMed](#)] [[Google Scholar](#)]
4. Gaston CM, Mitchell G: Information giving and decision-making in patients with advanced cancer: A systematic review. *Soc Sci Med* 2005;61:2252–2264 [[PubMed](#)] [[Google Scholar](#)]
5. Christakis NA, Lamont EB: Extent and determinants of error in doctors' prognoses in terminally ill patients: Prospective cohort study. *BMJ* 2000;320:469–472 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
6. Hagerty RG, Butow PN, Ellis PM, et al.: Communicating prognosis in cancer care: A systematic review of the literature. *Ann Oncol* 2005;16:1005–1053 [[PubMed](#)] [[Google Scholar](#)]
7. Downer J, Goldman R, Pinto R, et al.: The “surprise question” for predicting death in seriously ill patients: A systemic review and meta-analysis. *CMAJ* 2017;189:484–493 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
8. Larkin JR, Robinson MG, Bernacki RE, et al.: Estimating 1-year mortality for high-risk primary care patients using the “surprise” question. *JAMA Intern Med* 2016;176:1863–1865 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
9. Burke K, Coombes LH, Menezes A, Anderson AK: The “surprise question” in paediatric palliative care: A prospective cohort study. *Palliat Med* 2018;32:535–542 [[PubMed](#)] [[Google Scholar](#)]
10. Moss AH, Ganjoo J, Sharma S, et al.: Utility of the “surprise” question to identify dialysis patients with high mortality. *Clin J Am Soc Nephrol* 2008;3:1379–1384 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
11. Pang WF, Kwan BC, Chow KM, et al.: Predicting 12-month mortality for peritoneal dialysis patients using the “surprise” question. *Perit Dial Int* 2013;33:60–66 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

12. Javier AD, Figueroa R, Siew ED, et al.: Reliability and utility of the surprise question in CKD stages 4 to 5. *Am J Kidney Dis* 2017;70:93–101 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
13. Moss AH, Lunney JR, Culp S, et al.: Prognostic significance of the “surprise” question in cancer patients. *J Palliat Med* 2010;13:837–840 [[PubMed](#)] [[Google Scholar](#)]
14. Moroni M, Zocchi D, Bolognesi D, et al.: The “surprise” question in advanced cancer patients: A prospective study among general practitioners. *Palliat Med* 2014;28:959–964 [[PubMed](#)] [[Google Scholar](#)]
15. Lilley EJ, Gemunden SA, Kristo G, et al.: Utility of the “surprise” question in predicting survival among older patients with acute surgical conditions. *J Palliat Med* 2017;20:420–423 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
16. Ouchi K, Jambaulikar, George NR, et al.: The “surprise question” asked of emergency physicians may predict 12-month mortality among older emergency department patients. *J Pall Med* 2018;21:236–240 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
17. Rice J, Hunter L, Hsu AT, et al.: Using the “surprise question” in nursing homes: A prospective mixed-methods study. *J Palliat Care* 2018;33:9–18 [[PubMed](#)] [[Google Scholar](#)]
18. White N, Kupeli N, Vickerstaff V, Stone P: How accurate is the “surprise question” at identifying patients at the end of life? A systematic review and meta-analysis. *BMC Med* 2017;15:139. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
19. Romo RD, Lynn J: The utility and value of the “surprise question” for patients with serious illness. *CMAJ* 2017;189:E1072–E1073 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
20. Weissman DE, Meier DE: Identifying patients in need of a palliative care assessment in the hospital setting: A consensus report from the Center to Advance Palliative Care. *J Palliat Med* 2011;14:17–23 [[PubMed](#)] [[Google Scholar](#)]